

DAFTAR PUSTAKA

- Adenkambi AF, Ogunlesi TA, Olowu AO, Fetuga MB. Current trends in prevalence and aetiology of childhood congestive cardiac failure in Sagamu. *Journal of tropical pediatrics*. 2006;103-106.
- Andrews RE, Fenton MJ, Ridout DA, Burch M. New onset heart failure due to heart muscle disease in childhood. *Circulation*. 2008;79-84.
- Auerbach SR, Richmond ME, Lamour JM, Blume ED, Addonizio LJ, et al. BNP levels predict outcome in pediatric heart failure patients. *Circ heart fail*. 2010;3:606-11.
- Baghdady Y, Kamel Y, Elwan A. N-terminal pro-Brain natriuretic peptide in decompensated ventricular septal defect. *Arch Med Sci*. 2009;376-81.
- Beggs S, Thompson A, Nash R, Thompson A, Peterson G, editors. Cardiac failure in children. 17th Expert committee on the selection and the essential medical;2009;Geneva:TASP Press;2009.p.1-35.
- Bernus A, Wagner BD, Accurso F, Doran A, Kaess H, Ivy D. Brain natriuretic peptide levels in managing pediatric patients with pulmonary arterial hypertension. *CHEST*. 2009;135:745-51.
- Beteencourt PM. Clinical usefulness of B-type natriuretic peptide measurement: present and future perspectives. *Heart*. 2005;91:1489-94.
- Bonatto RS, Fioretto JR, Okoshi K, Matsubara BB, Padovani CR, et al. Percentiles curves of normal value of echocardiographic measurements in normal children from the central-southern region of the State of Sao Paulo, Brazil. *Depto de Pediatria: Faculdade de medicina de Botucatu UNESP*. 2005;651-60.
- Bradham WS, Bozkurt B, Gunasinghe H, Mann D, Spinale FG. Tumor necrosis factor alpha and myocardial remodelling in progression of heart failure: a current perspective. *Cardiovascular research*. 2002;53:822-30.
- Brunner M, Moeslinger T, Spieckerman PG. Echocardiography for teaching cardiac physiology in practical student courses. *Advan in Physiol Edu*. 1995;268:S1-9.
- Carleton PF. Fisiologi sistem kardiovaskuler. Dalam: Price SA, Wilson LM, editors. Patofisiologi konsep klinis proses-proses penyakit. Edisi 4. Jakarta;ECG. 1995. hal:480-95.
- Casals G, Ros J, Sionis A, Davidson MM, Ruiz MM, Jimenez W. Hypoxia induces B-type natriuretic peptide release in cell lines derived from human cardiomyocytes. *Am J Physiol Circ Physiol*. 2009;297:H550-55.

- Castello JM, Goodman DM, Green TP. A review of the natriuretic hormone system diagnostic and therapeutic potential in critically ill children. *Pediatr Crit Care Med*. 2006;7:308-16.
- Clerico A, Giannoni A, Vittorini S, Passino C. Thirty year of the heart as an endocrine organ: physiological role and clinical utility of cardiac natriuretic hormones. *Am J Physiol Heart Circ Physiol*. 2011;301:H12-20.
- Colak A, Cuhada S, Golcuk B, Ozogan O, Coker I. Effect of renal failure on N-terminal pro-brain natriuretic peptide in patients admitted to emergency department with acute dyspnea. *Anadolu Kardiyol Derg*. 2013;14:1-6.
- Crawford MH, Grant D, O'Rourke RA, Starling MR, Groves BM. Accuracy and reproducibility of new M-mode echocardiographic recommendations for measuring left ventricular dimensions. *Circulation*. 1980;61:137-43.
- Dahlan MS. Besar sampel dan cara pengambilan sampel. Jakarta. Salemba medika. 2010. hal. 76-7.
- Dan H, Yang L, Huixian T, Jinping G. Clinical value of plasma B-type natriuretic peptide assay in pediatric pneumonia accompanied by heart failure. *Experimental and therapeutic medicine*. 2015;2:175-9.
- Data rawat inap bangsal anak RSUP dr. M. Jamil tahun 2012 dan tahun 2013.
- De Bold, Bruneau, Kuroski ML. Mechanical and neuroendocrine regulation of the endocrine heart. *Cardiovascular research*. 1996;31:7-18.
- Ekure EN, Okoromah CA, Ajuluchukwu JN, Mbakwen A, Oladipo J. Diagnostic usefulness of N-terminal pro brain natriuretic peptide among children with heart failure in a tertiary hospital in Lagos, Nigeria. *WJG*. 2011;30:29-34.
- Emdin M, Vittorini S, Passino C, Clerico A. Old and new biomarkers of heart failure. *European Journal of heart failure*. 2009;11:331-35.
- Felker GM, Hasselbach V, Hernandez AF, O'Connor CM, Durham. Biomarker-guided therapy in chronic heart failure: A meta-analysis of randomized controlled trials. *American Heart Journal*. 2009;158:422-28.
- Foo R. Basic mechanism in apoptosis and heart failure. *Heart metab*. 2010;47:5-8.
- Francis GS, Sonnenblick EH, Tang WHW, Wilson PP. Pathophysiology of heart failure. In: Fuster V, O'Rourke R, Walsh RA, Wilson PP, King SB, Nash IS, et al, editors. *Hurst's the heart*. 2nd ed. New York: Mc Graw Hill medical; 2008. pp. 691- 12.

- Geva T, Vander velde ME. Imaging techniques: echocardiography, magnetic resonance imaging and computerized tomography. In; Keane JF, Lock JE, Flyer DC, editors. NADAS'S pediatric cardiology, 2rd ed. Philadelphia. Elsevier. 2006. pp. 145-82.
- Graca RL, Hasinger DC, Flynn PA, Sison CP, Nesin M, Auld P. Longitudinal changes of brain type natriuretic peptide in preterm neonatus. *Pediatrics*. 2006;117:2183-89.
- Haeling SV, Doehner W, Anker SD. Nutrition, metabolism, and the complex pathophysiology of cachexia in chronic heart failure. *Cardiovascular research*. 2007;73:298-09.
- Hsu DT, Pearson GD. Heart failure in children. Part I. History, etiology, and pathophysiology. *Circulation*. 2009;120:63-70.
- Ito T, Yoshimura M, Nakamura S, Nakayama M, Shimasaki Y, Harada E, et al. Inhibitory effect of natriuretic peptides on aldosterone synthase gene expression in cultured neonatal rat cardiocytes. *Circulation*. 2003;107:807-10.
- Katz AM, Konstam MA. The neurohormonal response in heart failure: Functional signaling. In: Katz AM, Konstam MA, editors. Heart failure. 2nd ed. Philadelphia: Lippincott Williams & Wilkins; 2009; pp. 87- 131.
- Kavga M, Varlaimis G, Giannopoulos A, Papadopolou KL, Varlaimis S, Bompotis G, et al. Correlation of plasma B-type natriuretic peptide with shunt volume in children with congenital heart disease involving left to right shunt. *Hellenic J Cardiol*. 2013; 192-98.
- Kim HN, Januzzi JL. Natriuretic peptide testing in heart failure. *Circulation*. 2011;123:2015-19.
- Koch A, Zink S, Singer H. B-type natriuretic peptide in paediatric patients with congenital heart disease. *European Heart Journal*. 2006;27:861-66.
- Konstam MA, Kramer DG, Patel AR, Maron MS, Udelson JE. Left ventricular remodelling in heart failure. *JACC*. 2011;4:98-108.
- Kumar R. Public health practice. Controlling rheumatic heart disease in developing countries. World health forum. 1995. Diunduh dari <http://whqlibdo.who.int> pada tanggal 22 november 2014.
- Lercher AH, Geiger R, Mair J, Url C, Tulzer G, Lechner E, et al. Utility of N-terminal pro-B-type natriuretic peptide to differentiate cardiac disease from noncardiac disease in young pediatric patients. *Clinical Chemistry*. 2006;52:1415-19.

- Lin CW, Zeng XL, Hu Jiang S, Wu T, Ping Wang J, Feng Shang J, et al. Role of NT-proBNP in the diagnosis of pediatric heart failure and investigation of novel combined diagnostic criteria. *Experimental and therapeutic medicine*. 2013;6:995-99.
- Lips DJ, deWindt LJ, van Kraaij DJW, Doevendans PA. Molecular determinants of myocardial hypertrophy and failure: alternative pathway for beneficial and maladaptive hypertrophy. *European heart journal*. 2003; 24:883-96.
- Macicek SM, Macias CG, Jefferies JL, Kim JJ, F Jack. Acute heart failure syndrome in the pediatric emergency department. *Pediatrics*. 2009;124:898-4.
- Madiyono B, Rahayuningsih SE, Sukardi R. Penanganan penyakit jantung pada bayi dan anak. *UKK kardiologi IDAI*. 2005: 55- 61.
- Maurellet JD, Liu PT. B-type natriuretic peptide in the management of heart failure. *Hong Kong Med J*. 2008;216-18.
- McGrath MR, Kuroski ML, de Bold AJ. The endocrine function of the heart. *TRENDS in Endocrinology and Metabolism*. 2005;16:469-75.
- McMurray JVJ. The role of angiotensin II receptor blockers in the management of heart failure. *Eurheartj*. 2005;7:J10-4.
- Mertens LL, Rigby ML, Horowitz ES, Anderson RH. Cross sectional echocardiographic and doppler imaging. In: Anderson RH, Baker EJ, Penny DJ, Redington AN, Rigby ML, Wernovsky G, editors. *Pediatric cardiology*. 3rd ed. Philadelphia. Elsevier. 2009. pp. 313-40.
- Mihl C, Dassen WRM, Kuipers H. Cardiac remodelling: concentric versus eccentric hypertrophy in strength and endurance athletes. *Netherlands Heart journal*. 2008;16:129-32.
- Mizuno Y, Yoshimura M, Yasue H, Sakamoto T, Ogawa H, Kugiyama K, et al. Aldosterone production is activated in failing ventricle in humans. *Circulation*. 2001;103:72-7.
- Moo GW. B-type natriuretic peptide in heart failure. *Curr Opin cardiol*. 2006;21:208-14.
- Morison LK, Harrison A, Krishnaswamy, Kazanegra R, Clopton P, Maisel A. Utility of B-type natriuretic peptide assay in differentiating congestive heart failure from lung disease in patients presenting with dyspnea. *J Am Coll Cardiol*. 2002;39:202-9.
- Nadas AS, Hauck AJ. Symposium on congestive heart failure: Pediatric aspect of congestive heart failure. *Circulation*. 1960;21:424-29.

- Nickenig G. Should angiotensin II receptor blockers and statins be combined? *Circulation*. 2004;110:1013-20.
- Nir A, Lindinger A, Rauh M, Bar-oz B, Laer S, Schwachtgen, Koch A, et al. NT-pro-B-type natriuretic peptide in infants and children: reference values based on combined data from four studies. *Pediatr Cardiol*. 2006;30:3-8.
- O'Byrne ML. Serum B-type natriuretic peptide as a biomarker for clinical severity of heart failure in pediatric heart disease. *Doris Duke Medical Student Journal*. 2005;5:53-9.
- Ogeng'O JA, Gatonga PM, Olabu BO, Nyamweya DK, Ongtera D. Pattern of congestive heart failure in Kenyan paediatric population. *CVJ Africa*. 2013;24:117-20.
- Oishi PE, Ridley AF, Datar SA, Fineman JR. B-type natriuretic peptide (BNP) in neonates, infants and children undergoing cardiac surgery. Front line of thoracic surgery. University of California. 2008. pp.199-08.
- Oyededeji OA, Oluwayemi IO, Oyededeji AT, Okeniyi JA, Fadero FF. Heart failure in Nigeria children. *The cardiology*. 2010:18-22.
- Park K Myung. *Pediatric Cardiology for practitioners*. 5th ed. Philadelphia: Elsevier; 2008:81-113; 461- 73.
- Price JF, Thomas AK, Grenier M, Eidem BW, Smith O, Denfield SW, et al. B-type natriuretic peptide predict adverse cardiovascular events in pediatric outpatients with chronic left ventricular systolic dysfunction. *Circulation*. 2006;1062-68.
- Rademaker MT, Espiner E. Hormones of cardiovascular system. In: Jameson JL, Kretser D, Grossman A, Marshall JC, Melmed S, De Groot L, et al, editors. *Endocrinology adult and pediatric*. 6th ed. Philadelphia. Elsevier;2010.pp. 2036-51.
- Rademaker MT, Espiner EA. Cardiovascular hormones. In: Besser GM, Thorner MO, editors. *Comprehensive clinical endocrinology*. 3rd ed. Philadelphia: Elsevier science;2002.pp. 486-99.
- Richards AM. The renin-angiotensin-aldosterone system and the cardiac natriuretic peptide. *Hearts*. 1996;76:36-44.
- Rusconi PG, Ludwig DA, Mas R, Hammon WG, Colan SD, Lipshutz SE. Serial measurement of serum NT-pro BNP as markers of left ventricle systolic dysfunction and remodelling in children with heart failure. *Am Heart J*. 2010;160:776-83. \

- Sastoasmoro S, Ismael SI. Dasar-dasar metodologi penelitian klinis. Edisi 4. Jakarta. Sagung seto. 2011. Hal.348-83.
- Sarangnga C. Korelasi negatif Fraksi Ejeksi Ventrikel Kiri dengan kadar N-terminal probrain natriuteric peptide pada anak dengan gagal jantung. Thesis. Universitas Udayana 2014.
- Seino Y, Ogawa A, Yamashita T, et al. Application of NT- proBNP and BNP measurements in cardiac care: a more discening marker for detection and evaluation af heart failure. *The european journal of heart failure*; 2004;295-300
- Srisawasdi P, Vanavanan S, Charoenpanichkit C, Kroll MH. The effect of renal dysfunction on BNP, NT-proBNP, and their ratio. *Am J Clin Pathol*. 2010;14-23.
- Sugimoto M, Manabe H, Nakau K, Furuya A, Okusima K, Fujiyasu H, et al. The role of NT-proBNP in the diagnosis of congestive heart failure. *Circulation journal*. 2010;74:998-5.
- Toora BD, Rajagopal G. Measurement of creatinine by Jaffe's reaction- Determination of sodium hydroxide required for maximum color development in standard, urine, protein free filtrate of serum. *Indian Journal of Experimental Biology*. 2002;352-54.
- Troughton RW, Richards M. B-natriuretic peptides and echocardiographic measures of cardiac structure and function. *J. Am. Coll. Cardiol. Img*. 2009;2:216-25.
- Van Empel VPM, Bertrand ATA, Hofstra L, Crijin HJ, Doevendans PA, Windt LJD. Myocyte apoptosis in heart failure. *Cardiovascular research*. 2005;67:21-9.
- Welish E, Kleesiek K, Haas N, Norozi K, Rauch R, Filler G. Aminoterminal pro B-type natriuretic peptide (NT-proBNP) levels for monitoring interventions in pediatric cardiac patients with stenotic lesions. *International Journal of Pediatrics*. 2009;1-6.
- Wu C, Wang Z hua, Li Z qin, Wang L feng. Effect of spironolactone on cardiac remodelling fter acute myocardial infarction. *World J Emerg Med*. 2013;4:48-53.
- Wu YR, Chen BS, Huang MR, Zhang YQ, Sun K, Chen S. N terminal pro brain natriuretic peptide in the diagnosis of congestive heart failure in pediatric patients with ventriculer septal defect. *World J Pediatr*. 2006;2:1-5.
- Zhou FJ, Zhou CY, Tian YJ, Xiao AJ, Li PL, Wang YH,et al. Diagnostic value of analysis of H-FABP, NT-proBNP, and CtnI in heart function in children with heart

disease and pneumonia. *European review for medical and pharmacological sciences*. 2014;18:1513-16.

Zoair AM, Mawlana WH, El-Bendari AS, Nada EA. Serum level of amino terminal of probrain natriuretic peptide as a diagnostic and prognostic bomarker in children with dilated cardiomyopathy. *Tanta Medical Journal*. 2014: 53-7.

